arXiv:q-alg/9702003v1 30 Jan 1997

Heisenberg Double Description of κ-Poincaré Algebra and κ-deformed Phase Space *

J. Lukierski[†] A. Nowicki[‡]

December 1996

Abstract

The κ-deformed dual pair of Poincaré algebra and Poincaré group is formulated in the framework of Heisenberg doubles. The covariant κ-deformed phase space is described in detail as a subalgebra. The realizations of proposed algebraic scheme are considered.

1 The Heisenberg double of κ-Poincaré algebra

It is known that from a pair of dual Hopf algebras describing the quantum symmetry group $\mathcal{A}$ and its dual quantum Lie algebra $\mathcal{A}^*$ one can construct three different double algebras: Drinfeld double and Drinfeld codouble, both with Hopf algebra structure, and Heisenberg double, which is not a Hopf algebra. Heisenberg double describes the semidirect product of the algebra of vector fields acting on the functions of quantum group and represents algebraic generalization of the notion of cotangent double on the group, i.e. describes the generalization of the phase space for the group manifolds.

In the present note we apply the notion of Heisenberg double of a bialgebra (see [1]) in the case of the κ-deformation of Poincaré algebra and its dual group [2, 3]. The Heisenberg double of κ-Poincaré algebra $\mathcal{H}(\mathcal{P}_\kappa)$ contains as its subalgebra another Heisenberg double which turns out to be the κ-deformed phase space.
We discuss in κ-deformed phase space the deformed uncertainty relations for this phase space which depend on two parameters: the Planck's constant $\hbar$ and the deformation parameter κ with mass dimension. One of the important problems in the phase space-formalism is the canonical change of phase space coordinates, leaving invariant the Heisenberg canonical commutation relations. In the Heisenberg double description the

---

[*] Supported by KBN grant 2P30208706

[†] Institute for Theoretical Physics, University of Wrocław, pl. Maxa Borna 9, 50-204 Wrocław, Poland.

[‡] Institute of Physics, Pedagogical University, pl. Słowiański 6, 65-029 Zielona Góra, Poland.

object which we would like to leave invariant under canonical transformations are the scalar products defining the duality relations. The problem of changing the basis in dual configuration space introduced by the nonlinear changes of fourmomentum generators we formulate in the language of the finite difference equations.

The deformation of the Heisenberg commutation relations has been studied in general algebraic framework by Kempf [4]. The deformation of classical relativistic symplectic structure for κ-Poincaré was discussed earlier in [5, 6]. Recently the link between the noncommutativity of the spacetime coordinates in κ-deformed Minkowski space and the effects of quantum gravity was considered also by Amelino-Camelia [7].

1.1 Heisenberg double

Let $\{A_a\}$ be a linear basis of bialgebra $\mathcal{A}$ with the following

- *multiplication relations*:

$$A_a \circ A_b = f_{ab}^c A_c \tag{1.1a}$$

- *co-multiplication relations*:

$$\Delta(A_a) = h_a^{bc} A_b \otimes A_c \tag{1.1b}$$

We recall the following definition [1]

Def. *The Heisenberg double $\mathcal{H}(\mathcal{A})$ of bialgebra $\mathcal{A}$ is an associative algebra with a basis $\{A_a, B^b\}$ satisfying*

$$A_a \circ A_b = f_{ab}^c A_c \tag{1.2a}$$

$$B^a \circ B^b = h_c^{ab} B^c \tag{1.2b}$$

$$B^a \circ A_b = f_{ed}^a h_b^{ce} A_c \circ B^d \tag{1.2c}$$

The bialgebra $\mathcal{A}$ and dual $\mathcal{A}^*$ are equivalent to subalgebras spanned by the basis $\{A_a\}$ and $\{B^b\}$ respectively.

The multiplication given by the formula (1.2c) can be derived from a left action of a dual algebra $\mathcal{A}^*$ on $\mathcal{A}$ defined as follows

$$b \triangleright a = a_{(1)} < b, a_{(2)} > \qquad\qquad b \in \mathcal{A}^*, a \in \mathcal{A} \tag{1.2d}$$

where we use Sweedler's notation for coproduct $\Delta(a) = a_{(1)} \otimes a_{(2)}$. In particular using duality relation between multiplication and comultiplication one gets

$$b \triangleright a\tilde{a} = (a\tilde{a})_{(1)} < b, (a\tilde{a})_{(2)} >= (a\tilde{a})_{(1)} < \Delta(b), a_{(2)} \otimes \tilde{a}_{(2)} >=$$
$$= (a\tilde{a})_{(1)} < b_{(1)}, a_{(2)} >< b_{(2)}, \tilde{a}_{(2)} >= a_{(1)} < b_{(1)}, a_{(2)} > (\tilde{a}_{(1)} < b_{(2)}, \tilde{a}_{(2)} >) =$$
$$= a_{(1)} < b_{(1)}, a_{(2)} > (b_{(2)} \triangleright \tilde{a}) = (b \circ a) \triangleright \tilde{a}$$

and we obtain

$$b \circ a = a_{(1)} < b_{(1)}, a_{(2)} > b_{(2)} \tag{1.2e}$$

The multiplication (1.2c) is equivalent to the commutation relations (1.2e). In particular, for a linear basis $\{A_a\}$ and $\{B^b\}$ we have

$$\Delta(A_a) = h_a^{ce} A_c \otimes A_e = A_{a(1)} \otimes A_{a(2)}$$

$$\Delta(B^b) = f_{cd}^{b} B^c \otimes B^d = B^b_{(1)} \otimes B^b_{(2)}$$

and the formula (1.2e) becomes (1.2c).

1.2 κ-Poincaré algebra

We choose the following realization of the κ-Poincaré algebra in bicrossproduct basis [2, 8]
($\mu, \nu, \lambda, \sigma = 0, 1, 2, 3; i, j = 1, 2, 3$ and $g^{\mu\nu} = g_{\mu\nu} = (-1, 1, 1, 1)$)

-*algebra sector*:

$$\begin{aligned}
&[P_\mu, P_\nu] = 0 \\
&[M_{\mu\nu}, M_{\lambda\sigma}] = i\hbar \left(g_{\mu\sigma} M_{\nu\lambda} + g_{\nu\lambda} M_{\mu\sigma} - g_{\mu\lambda} M_{\nu\sigma} - g_{\nu\sigma} M_{\mu\lambda} \right) \\
&[M_{ij}, P_\mu] = -i\hbar \left(g_{i\mu} P_j - g_{j\mu} P_i \right) \\
&[M_{i0}, P_0] = i\hbar P_i \\
&[M_{i0}, P_j] = i\delta_{ij} \left(\hbar^2 \kappa \sinh(\tfrac{P_0}{\hbar\kappa}) e^{-\frac{P_0}{\hbar\kappa}} + \tfrac{1}{2\kappa} \vec{P}^2 \right) - \tfrac{i}{\kappa} P_i P_j
\end{aligned} \tag{1.3}$$

-*coalgebra sector*:

$$\begin{aligned}
\Delta(M_{ij}) &= M_{ij} \otimes I + I \otimes M_{ij} \\
\Delta(M_{k0}) &= M_{k0} \otimes e^{-\frac{P_0}{\hbar\kappa}} + I \otimes M_{k0} + \tfrac{1}{\hbar\kappa} M_{kl} \otimes P_l \\
\Delta(P_0) &= P_0 \otimes I + I \otimes P_0 \\
\Delta(P_k) &= P_k \otimes e^{-\frac{P_0}{\hbar\kappa}} + I \otimes P_k
\end{aligned} \tag{1.4}$$

where $\hbar$ denotes Plank's constant and κ deformation parameter. The formulas for the antipode and counit are omitted because they are not essential for this construction.

1.3 κ-Poincaré group

Using the following duality relations

$$< x^\mu, P_\nu > = \ i\hbar \delta^\mu_\nu \qquad < \Lambda^\mu{}_\nu, M_{\alpha\beta} > = \ i\hbar (\delta^\mu_\alpha g_{\nu\beta} \ - \ \delta^\mu_\beta g_{\nu\alpha}) \tag{1.5}$$

we obtain the commutation relations defining κ-Poincaré group [3, 8] in the form

-*algebra sector*:

$$\begin{aligned}
[x^\mu, x^\nu] &= \tfrac{i}{\kappa}(\delta_0^\mu x^\nu - \delta_0^\nu x^\mu) \\
[\Lambda_\nu^\mu, x^\lambda] &= -\tfrac{i}{\kappa}\left((\Lambda_0^\mu - \delta_0^\mu)\Lambda_\nu^\lambda + (\Lambda_\nu^0 - \delta_\nu^0)g^{\mu\lambda}\right) \\
[\Lambda_\nu^\mu, \Lambda_\beta^\alpha] &= 0 \qquad (1.6)
\end{aligned}$$

-*coalgebra sector*:

$$\begin{aligned}
\Delta(x^\mu) &= \Lambda^\mu{}_\alpha \otimes x^\alpha \ + \ x^\mu \otimes I \\
\Delta(\Lambda^\mu{}_\nu) &= \Lambda^\mu{}_\alpha \otimes \Lambda^\alpha{}_\nu \qquad (1.7)
\end{aligned}$$

The commutation relations (1.3) and (1.6) one can supplement by the following relations obtained from (1.2c) or (1.2*e*)

-*cross relations*:

$$\begin{aligned}
[P_k, x_l] &= -i\hbar\delta_{kl} & [P_0, x_0] &= \imath\hbar \\
[P_k, x_0] &= -\tfrac{i}{\kappa}P_k & [P_0, x_l] &= 0 \\
[P_\mu, \Lambda_\beta^\alpha] &= 0 \\
[M_{\alpha\beta}, \Lambda_\nu^\mu] &= i\hbar\left(\delta_\beta^\mu\Lambda_{\alpha\nu} - \delta_\alpha^\mu\Lambda_{\beta\nu}\right) \\
[M_{\alpha\beta}, x^\mu] &= i\hbar\left(\delta_\beta^\mu x_\alpha - \delta_\alpha^\mu x_\beta\right) + \tfrac{i}{\kappa}\left(\delta_\beta^0 M_\alpha{}^\mu - \delta_\alpha^0 M_\beta{}^\mu\right) \qquad (1.8)
\end{aligned}$$

where

$$M_\alpha{}^\mu \ = \ g^{\mu\rho}M_{\alpha\rho} \qquad M^\mu{}_\alpha \ = \ g^{\mu\rho}M_{\rho\alpha} \qquad (1.9)$$

The relations (1.3), (1.6) and (1.8) give us the Heisenberg double $\mathcal{H}(\mathcal{P}_\kappa)$ of κ-Poincaré algebra in terms of commutation relations.

2 κ-Deformed Phase Space

Let us consider subalgebra of $\mathcal{H}(\mathcal{P}_\kappa)$ given by the following

- *commutation relations*:

$$\begin{aligned}
[x_k, x_l] &= 0 & [P_\mu, P_\nu] &= 0 \\
[x_0, x_k] &= \tfrac{i}{\kappa}x_k \\
[x_k, P_l] &= i\hbar\delta_{kl} & [x_k, P_0] &= 0 \\
[x_0, P_l] &= \tfrac{i}{\kappa}P_l & [x_0, P_0] &= -i\hbar
\end{aligned} \qquad (2.1)$$

which describe the κ-deformed phase space. For $\kappa \to \infty$ we get the standard nondeformed phase space satisfying the Heisenberg commutation relations. This phase space one can obtain immediately using the relations (1.2) to the commuting fourmomentum algebra with noncocommutative coproduct (1.4).

2.1 Realization of κ-deformed phase space algebra in terms of standard phase space variables

It is easy to see that the commutation relations (2.1) can be realized in a Hilbert space in terms of standard undeformed momentum and position operators satisfying the Heisenberg commutation relations [6]

$$[\hat{x}_\mu, \hat{p}_\nu] \;=\; i\hbar g_{\mu\nu} \tag{2.2}$$

then we can define:

$$x_k = \hat{x}_k \quad x_0 = \hat{x}_0 + \tfrac{1}{2\kappa\hbar}(\hat{x}_i\hat{p}_i + \hat{p}_i\hat{x}_i) \quad P_\mu = \hat{p}_\mu \tag{2.3}$$

Equivalently, we can describe this transformation introducing the operator

$$U \;=\; e^{\frac{i}{2\kappa\hbar^2}(\hat{x}_k\hat{p}_k + \hat{p}_k\hat{x}_k)\hat{p}_0} \tag{2.5}$$

then we get

$$x_k \;=\; \hat{x}_k \qquad x_0 \;=\; U\hat{x}_0 U^{-1} \qquad P_\mu \;=\; \hat{p}_\mu \tag{2.6}$$

It appears that for an arbitrary invertible operator U , the algebra generated by x_μ, P_ν satisfies the Jacobi identity. Therefore, such an operator U describes some general class of deformations of the standard phase space given by (2.2)

2.2 κ-deformed Heisenberg uncertainty relations

Introducing the dispersion of the observable a in quantum mechanical sense by:

$$\Delta(a) \;=\; \sqrt{< a^2 > - < a >^2} \tag{2.7}$$

we obtain the nonvanishing κ-deformed uncertainty relations in κ-deformed phase space as form

$$\begin{aligned}
&\Delta(x_0)\Delta(x_k) \geq \tfrac{1}{2\kappa}| < x_k > |\\
&\Delta(P_k)\Delta(x_l) \geq \tfrac{1}{2}\hbar\delta_{kl}\\
&\Delta(P_0)\Delta(x_0) \geq \tfrac{1}{2}\hbar\\
&\Delta(P_k)\Delta(x_0) \geq \tfrac{1}{2\kappa}| < P_k > |
\end{aligned} \tag{2.8}$$

which become the standard ones in the limit $\kappa \to \infty$.

2.3 Remark on changing the basis

It is obvious, that for our choice of the basis in phase space, the following general duality relations hold ($r, s = 1, 2, 3$):

$$< x_0^k x_s^l, P_r^n P_0^m >= (\hbar)^{k+l}\delta_{rs}\delta^{ln}\delta^{km}k!l!(-i)^k(i)^l \tag{2.9a}$$

$$< x_s^l x_0^k, P_r^n P_0^m >= \left\{ \begin{array}{ll} (\hbar)^{k+l}\delta_{rs}\delta^{ln}\frac{k!l!}{(k-m)!}\left(-\frac{n}{\kappa\hbar}\right)^{k-m}(-i)^k(i)^l & k\geq m \\ 0 & k<m \end{array} \right\} \tag{2.9b}$$

and for any polynomial functions $\psi(x_0,\vec{x})$ and $f(P_0,\vec{P})$ the duality pairing has the form

$$<: \psi(x_0,\vec{x}) :, f(P_0,\vec{P}) >= f(-i\hbar\partial_0, i\hbar\vec{\bigtriangledown})\psi(x_0,\vec{x})|_{x=0} \tag{2.10}$$

where $: \psi :$ denotes odered function ψ with all powers of x_0 to the left.
Let us consider the following change of the basis of κ-Poincaré algebra:

$$\tilde{P}_i = P_i e^{\frac{P_0}{\kappa\hbar}} = P_i k(P_0) \tag{2.11a}$$

$$\tilde{P}_0 = \kappa\hbar\sinh\left(\frac{P_0}{\kappa\hbar}\right) + \frac{1}{2\kappa\hbar}\vec{P}^2 k(P_0) = h(P_0,\vec{P}^2) \tag{2.11b}$$

which transforms to the standard, nondeformed basis of the Poincaré algebra.

From the duality relations for the phase space, the corresponding space-time $\{\tilde{x}_l, \tilde{x}_0\}$ related to the transformed momentum space $\{\tilde{P}_l, \tilde{P}_0\}$ i.e.

$$< \tilde{x}_l, \tilde{P}_k >= i\hbar\delta_{kl} \qquad < \tilde{x}_0, \tilde{P}_0 >= -i\hbar \tag{2.12}$$

is given by

$$\tilde{x}_l =: F_l(x_0,\vec{x}) : \qquad \tilde{x}_0 =: F_0(x_0,\vec{x}) : \tag{2.13}$$

where the functions F_l and F_0 satisfy the following differential equations:

$$\left(\frac{\kappa}{2}(1-e^{\frac{2i}{\kappa}\partial_0}) - \frac{1}{2\kappa}\triangle\right)^m \partial_r^n \left(F_0^k(x_0 - i\frac{m+n}{\kappa},\vec{x})\cdot\right. \tag{2.14}$$

$$\left. F_s^l(x_0 - i\frac{m+n}{\kappa},\vec{x})\right) |_{x=0} = (-i)^k k! l! \delta_{rs}\delta^{ln}\delta^{km} \tag{2.15}$$

in particular for $l = n = 0$

$$\left(\frac{\kappa}{2}(1-e^{\frac{2i}{\kappa}\partial_0}) - \frac{1}{2\kappa}\triangle\right)^m F_0^k(x_0 - i\frac{m}{\kappa},\vec{x}) |_{x=0} = (-i)^k k! \delta^{km} \tag{2.16a}$$

for $k = m = 0$

$$\partial_r^n F_s^l(-i\frac{n}{\kappa},\vec{x}) |_{\vec{x}=0} = n! \delta_{rs}\delta^{ln} \tag{2.16b}$$

2.4 Final remarks

In this note we presented the κ-deformed phase space for the standard version of κ-deformation, which introduces the "quantized" nature of time coordinate. An analogous discussion can be presented in the case of generalized κ-deformation (see [9, 10] and the lecture of Maslanka) describing fourdimensional space-time relativistic symmetries with one arbitrary direction becoming "quantum" after the κ-deformation. It should be recalled that the clasification of the deformations of D=4 Poincaré group has been presented recently by Podles and Woronowicz [11]. Taking into account the clasification of classical r-matrices or D=4 Poincaré algebra by Zakrzewski [10] it would be interesting to describe all possible deformed quantum phase spaces obtained by the Heisenberg double construction for known variety of deformed D=4 fourdimensional relativistic symmetries.

Acknowledgments

The authors would like to thank the organizers of the conference, in particular prof. V.D. Doebner and prof. V.K. Dobrev, for their hospitality in Goslar.

References

[1] N.Yu. Reshetikhin, M.A. Semenov-Tian-Shansky, *Lett. Math. Phys* **19** (1990) 133.
A.Yu. Alekseev, L.D. Faddeev, *Commun. Math. Phys.* **141** (1991) 413.
R.M. Kashaev, "Heisenberg double and pentagon relation", preprint ENSLAPP-L- 512/95.

[2] J. Lukierski, A. Nowicki, H. Ruegg, *Phys. Lett.* **B 293** (1993) 419.

[3] S. Zakrzewski, *Journ. Phys.* **A 27** (1994) 2075.

[4] A. Kempf, Journ. Math. Phys. **35**, 4483 (1994); hep-th/9602085

[5] S. Zakrzewski, in "Quantum groups: Formalism and Applications", Proceedings of XXX-th Karpacz Winter School, Polish Scientific. Publ., Warsaw 1995, p. 573.

[6] J. Lukierski, H. Ruegg, W.J. Zakrzewski, *Ann. of Phys.* **243** (1995) 90.

[7] G. Amelino-Camelia, Oxford Univ. preprint OUTP-96-54P, gr-qc/9611016, Phys. Lett. **B**, in press.

[8] P. Kosiński, P. Maślanka, "The duality between κ-Poincaré algebra and κ-Poincaré group", Lodz preprint IM UL 3/1994.

[9] P. Kosiński, P. Maślanka, in: *From field theory to quantum groups*, eds. B. Jancewicz and J. Sobczyk, World Scientific 1996, p. 41.

[10] S. Zakrzewski, in: *Quantum Groups: Formalism and Applications*, Proceedings of XXXth Karpacz Winter School, Polish Scientific Publ., Warsaw 1995, p.433.; see also q-alg 9602001.

[11] P. Podleś, S.L. Woronowicz, *Comm. Math. Phys.* **178**, (1996) 61.